

Microsatellite markers in analysis of resistance to coffee leaf miner in Arabica coffee

Gabriella Santos Pereira⁽¹⁾, Lilian Padilha⁽²⁾, Edila Vilela Resende Von Pinho⁽¹⁾,
Rita de Kássia Siqueira Teixeira⁽¹⁾, Carlos Henrique Siqueira de Carvalho⁽²⁾, Mirian Peres Maluf⁽²⁾
and Bruna Line de Carvalho⁽¹⁾

⁽¹⁾Universidade Federal de Lavras, Caixa Postal 3037, CEP 37200-000 Lavras, MG, Brazil. E-mail: gabipereira87@yahoo.com.br, edila@dag.ufla.br, ritaaadekassia@hotmail.com, brunalcarvalho@hotmail.com ⁽²⁾Embrapa Café, Parque Estação Biológica, s/nº, CEP 70770-901 Brasília, DF, Brazil. E-mail: lilian.padilha@embrapa.br, carlos.carvalho@embrapa.br, mirian.maluf@embrapa.br

Abstract – The objective of this work was to analyze coffee (*Coffea arabica*) genotypes resistant to the coffee leaf miner (*Leucoptera coffeella*) using microsatellite markers. Sixty-six loci were evaluated, of which 63 were obtained from the Brazilian Coffee Expressed Sequence Tag (EST) database. These loci were amplified in bulks of individuals from F₅ progenies of 'Siriema' (*C. arabica* x *C. racemosa*) resistant and susceptible to the insect, in eight samples of *C. racemosa*, and in a F₆ population of 'Siriema' with 91 individuals segregating for resistance to the leaf miner. Polymorphisms were verified for two simple sequence repeat (SSR) loci in bulks of the susceptible progenies. The two polymorphic alleles were present in around 70% of the susceptible genotypes in F₅ and in approximately 90% of the susceptible individuals in F₆. However, the polymorphic EST-SSR markers among populations contrasting for resistance to leaf miner were not correlated to the evaluated characteristics. SSR markers show inter- and intraspecific polymorphism in *C. arabica* and *C. racemosa*.

Index terms: *Coffea arabica*, *Leucoptera coffeella*, EST-SSR, genotypes.

Marcadores microssatélites na análise de resistência ao bicho-mineiro em café arábica

Resumo – O objetivo deste trabalho foi analisar genótipos de cafeeiro (*Coffea arabica*) resistentes ao bicho-mineiro (*Leucoptera coffeella*), com uso de marcadores microssatélites. Foram avaliados 66 locos, dos quais 63 foram obtidos do banco de sequências expressas (EST) de café. Esses locos foram amplificados em “bulks” de progênies F₅ de 'Siriema' (*C. arabica* x *C. racemosa*) resistentes e suscetíveis ao ataque da praga, em oito acessos de *C. racemosa*, e em uma população F₆ de 'Siriema' com 91 indivíduos que segregam para resistência ao bicho-mineiro. Os polimorfismos foram verificados em dois locos de marcadores microssatélites (SSR), nos “bulks” das progênies suscetíveis. Os dois alelos polimórficos estiveram presentes em torno de 70% dos genótipos suscetíveis em F₅ e em, aproximadamente, 90% dos indivíduos suscetíveis em F₆. Entretanto, os marcadores EST-SSR polimórficos entre populações contrastantes para a resistência ao bicho-mineiro não foram correlacionados às características avaliadas. Os marcadores SSR apresentam polimorfismo inter e intraespecífico em *C. arabica* e *C. racemosa*.

Termos para indexação: *Coffea arabica*, *Leucoptera coffeella*, EST-SSR, genótipos.

Introduction

The coffee leaf miner [*Leucoptera coffeella* (Guérin-Mèneville)] is considered the main pest in Brazilian coffee plantations. This microlepidoptera reduces the photosynthetic area of leaves, causing large losses in coffee production (Souza et al., 1998). Up to the present, commercial cultivars of Arabica coffee (*Coffea arabica* L.) registered in Brazil are susceptible to the coffee leaf miner, making chemical control of this pest indispensable. The use of resistant cultivars is an advantageous alternative for the control of biotic stresses. This strategy promotes a greater

competitiveness for farmers, since it allows greater yields by reducing losses caused by the attack of pests and diseases, production costs and pollution of the environment by the application of agricultural chemicals.

The *Coffea* genus has 103 species (Davis et al., 2006) with 2n=22 chromosomes. The exception is *C. arabica*, an allotetraploid, with 44 chromosomes, of which 22 came from *C. eugenioides* S. Moore and the rest from *C. canephora* Pierre ex A. Froehner (Lashermes et al., 1999). *C. arabica* has a narrow genetic base due to the small quantity of seeds originally introduced in Brazil associated with the autogamous nature of

this species. In addition, the predominance of mass selection and pedigree methods in breeding strategies has also contributed to reduce genetic diversity in *C. arabica*.

The resistance to *L. coffeella* in *Coffea* spp. was identified in the Instituto Agronômico (IAC) coffee breeding program in a fertile plant derived from a natural cross between *C. racemosa* Ruiz & Pav. and the Blue Mountain cultivar of *C. arabica*. Hybrid populations obtained from the combination of backcrossing and pedigree methods are established in the field, but these populations are still segregating for this trait (Guerreiro Filho, 2006).

The leaf miner resistance introduced from *C. racemosa* is controlled by two dominant complementary genes, Lm1 and Lm2 (Guerreiro Filho et al., 1999). However, the difficulty in stabilizing this trait in the genotypes indicates the need for more studies. So far, molecular studies of leaf miner resistance did not help to determine the nature of this resistance in Arabica coffee plants (Mondego et al., 2005; Pinto et al., 2007).

Microsatellite or simple sequence repeat (SSR) markers are based on repetitive DNA sequences and occur in the genome of all organisms. These markers are highly variable, and multiallelic genetic loci stand out for codominance and high content of polymorphism information (Tautz & Renz, 1984; Gupta & Varshney, 2000; Morgante et al., 2002). Microsatellites can be scored by many methods, which include derivation from enriched genomic libraries and screening of cDNA libraries from public databases, such as GenBank, from related species and from EST (expressed sequence tag) databases. EST-SSR have some intrinsic advantages in that they are quick to elucidate – by electronic sorting –, abundant, unbiased in repeat type and present in gene-rich areas (Scott, 2001). The Brazilian Coffee EST database has more than 200 thousand DNA sequences (Vieira et al., 2006).

The objective of this work was to analyze *C. arabica* genotypes resistant to the coffee leaf miner (*Leucoptera coffeella*) using microsatellite markers.

Materials and Methods

The genetic materials were obtained from the experimental fields of Coromandel and Boa Esperança, MG, Brazil, of the Fundação Procafé and from the Centro de Café of Instituto Agronômico (IAC),

Campinas, SP, Brazil. Molecular analyses were done at the Laboratório de Análise de Sementes e Biotecnologia of the Universidade Federal de Lavras, Lavras, MG, Brazil.

Leaves from the following genetic materials were used: eight *C. racemosa* accessions from the IAC coffee germplasm bank, which were identified as CR1, CR2, CR3, CR4, CR6, CR8, CR9 and CR10; 'Siriema' progenies from the Coromandel coffee field, of which ten were resistant F₅ progenies (R3, R23, R29, R30, R33, R34, R37, R43, R46 and R49) and ten were susceptible F₅ progenies (S1, S4, S8, S13, S15, S21, S24, S38, S42 and S48) – each progeny was composed of ten plants –; and 91 F₆ 'Siriema' plants from Boa Esperança, MG, Brazil. This population was used to evaluate segregating loci that were detected in F₅ progenies.

After the *C. arabica* x *C. racemosa* cross, the material passed through two backcrossings with *C. arabica* at IAC, and the segregant IAC 1195-5-6-2 was selected. C1195-5-6-2-119, with high resistance to coffee leaf miner, was selected by the Procafé breeding program and crossed with 'Catimor UFV 417' to incorporate rust resistance. From this cross, segregant 842 was selected, with resistance to rust and to leaf miner. Plants 842-1 and 842-4, designated 'Siriema', were advanced in their generations by the pedigree method, and currently are in the F₅ generation (Carvalho et al., 2008).

The 'Siriema' F₅ progenies were in the field for five years under evaluation for tolerance to the coffee leaf miner. The F₆ population was characterized at the IAC greenhouse (Guerreiro Filho et al., 1999). Lesion reaction was observed in leaf disks infested by *L. coffeella*, according to four classes: resistant, plants with point lesions; moderately resistant, plants with small threadlike lesions; moderately susceptible, plants with large threadlike lesions; and susceptible, plants with large irregular lesions.

Thirty-five EST-SSRs di- and trinucleotides, already available in the EST Coffee Genome database, were selected as a strategy to increase the possibility of obtaining polymorphic markers. These EST-SSR are more polymorphic than tetra-, penta- or hexanucleotides (Poncet et al., 2004). Information about the library, repetition patterns and the frequency of repetitions were extracted from the database (Table 1). Parameters for primer design considered 54 to 65°C melting

temperature (Tm) and 40 to 50% CG bases; amplicons ranged from 100 to 300 nucleotides; and primers had 18 to 22 bases (Table 2).

In addition, 28 EST-SSR primers (Pinto et al., 2007) and three more SSR loci, already characterized

for coffee biotic stress (Herrera et al., 2009), were used: Sat 225 (F: 5'CAT GCC ATC ATC AAT TCC AT3'; R: 5'TTA CTG CTC ATC ATT CCG CA3'), Sat 229 (F: 5'TTC TAA GTT GTT AAA CGA GAC GCT TA3'; R: 5'TTC CTC CAT GCC CAT ATT G3')

Table 1. EST-SSR loci selected from the EST Coffee Genome database.

Locus	Genomic library ⁽¹⁾	Motif	Repetition frequency	Expected size (bp)	Observed size (bp)	Number of alleles
LEG1	CL2, PA1	GA	21.5	222	180–200; 600	3
LEG2	AR1	AC	15.5	226	180–220	2
LEG3	FB1, FR2	AT	26.0	379	200	3
LEG4	AR1, LV9, RX1	AT	26.0	250	– ⁽²⁾	–
LEG5	LV9, LP1, SS1	AT	80.0	599	210–500	4
LEG6	CL2	TA	15.0	200	–	–
LEG7	RT8, CS1, RX1	TC	15.5	237	–	–
LEG8	FB1, LP1	GAG	16.7	218	–	–
LEG9 ⁽³⁾	FB4	GAT	19.0	235	120–230	4
LEG10	LV9	AG	15.0	228	230	1
LEG11 ⁽³⁾	BP1, CB1, EA1, EB1, EM1, FR4, IC1, RX1, S13, RM1	TGG	27.3	281	150–200	2
LEG12 ⁽³⁾	BP1, LV5, S13	TC	20.5	223	150–210	3
LEG13 ⁽³⁾	EM1, S13, SH2	GAA	18.0	249	150–300	3
LEG14	RX1, FR1	AT	20.5	210	150–180	2
LEG15	CL2	CT	16.5	216	200–350	4
LEG16	CL2, FB2, LV5, RT5, RX1	CA	27.5	244	–	–
LEG17	FB4, PC1	CT	22.0	190	130–180	2
LEG18	CL2, EA1, FR1, S13, RX1	GCT	10.7	249	250–300	2
LEG19	CB1, CS1, FB1, FR2, IA2, LV5, LV9, RT5, RX1	GAA	22.0	291	150–350	4
LEG20	CS1, FB1, LV5, RX1, RM1	TA	22.0	282	180–300	2
LEG21	EA1, S13, SH2	AGG	14.0	290	160–290	2
LEG22 ⁽³⁾	EA1	TC	21.5	286	300	1
LEG23	CA1, RT8, FR4, EA1, RX1, S13	GA	23.0	218	–	–
LEG24	CA1, FR2, FB1, FB2, BP1, CB1, EA1, IA2, LP1, PA1, S13, SH3, FV2	TC	27.0	176	300	1
LEG25	LV5, CL2, CB1, IA2	AT	17.5	183	270–300	2
LEG26 ⁽³⁾	CS1	ATA	12.7	197	190–230	2
LEG27	FB1, CB1	TA	20.0	231	180–300	3
LEG28 ⁽³⁾	BP1, FR4, RT3	AGT	11.0	250	250–280	3
LEG29	CA1, CL2, CS1, EA1, FR2, FR4, S13, SH2	AG	16.0	163	170–200	2
LEG30	LV5	CTG	19.0	239	240–320	2
LEG31	AR1, CS1, LV5, S13	CCT	13.3	247	250	1
LEG32 ⁽³⁾	CS1, RX1, S13	AGA	19.0	214	210–300	3
LEG33	CS1, RT8	AG	56.0	219	–	–
LEG34	CS1, FR4	TA	25.5	212	200–280	3
LEG35	BP1, CB1, CS1, FR4, IC1, RX1, S13, RM1	AT	23.5	210	–	–

⁽¹⁾CA1, non embryogenic callus; LV5, pointer leaf – orthotropic branch without Bion – short; LV9, pointer leaf – plagiotropic branches without Bion – short; RT3, root without Bion; RT5, root with Bion; CL2, hypocotyl callus with and without Bion; FR1, flower bud no. 6, lead no. 1, fruit stage 1 and 2 – long; FR2, flower bud no. 6, lead no. 1, fruit stage 1 and 2 – short; FB1, flower bud stage 1 and 2 – long; FB2, flower bud stage 1 and 2 – short; FB4, flower bud stage 3 and 4; RT8, root and cells in suspension with aluminum; BP1, cells in suspension, root and leaf with Bion; RM1, leaf infested with coffee leaf miner and rust; CB1, cells in suspension with Bion and brassinosteroids; CS1, cells in suspension with mannose, NaCl and KCl; FR4, racemosa fruits; EA1, embryogenic callus of *Coffea arabica*; EB1, zygotic embryo (immature fruits); IC1, non embryogenic progeny (*C. arabica* leaf) without induction of 2.4D; PC1, non embryogenic progeny (*C. arabica* leaf) with induction of 2.4D; IA2, embryogenic progeny (*C. arabica* leaf) with induction of 2.4D; AR1, leaves treated with arachidonic acid; LP1, seedlings treated with arachidonic acid; RX1, branches infected with *Xylella* spp.; PA1, embryogenic progeny (*C. arabica* primary callus); S13, seeds at the beginning of germination; SH2, water stress in the field (tissue pool – leaves, branches, gemmae, etc.); SH3, water stress in the field (*C. canephora* clone tolerant to drought); FV2, leaf stage 1, 2 and 3 (*C. racemosa*); SS1, irrigated in the field (tissue pool – leaves, branches, gemmae, etc.); EM1, embryo (seed). ⁽²⁾Não houve amplificação. ⁽³⁾Polymorphic loci.

and Sat 259 (F: 5'GCC AAT TGT GCA AAG TGC T-3'; R: 5'ATT CAT GGG GCC TTT GTC TT3').

All the primers were screened in the *C. racemosa* and F_5 progenies. The polymorphic oligonucleotides were used to amplify DNA fragments of each individual of the F_5 progenies and of the F_6 individuals of the population. Genomic DNA was extracted according to Ferreira & Grattapaglia (1996) by the CTAB 2% method. Final reaction conditions were 1X PCR buffer (Tris-HCl, 200 $\mu\text{mol mL}^{-1}$; pH 8.4; KCl, 500 $\mu\text{mol mL}^{-1}$; 2 $\mu\text{mol mL}^{-1}$ of MgCl_2 ; 400 nmol mL^{-1} of dNTPs; 0.3 nmol mL^{-1} of each primer; 0.5 U of DNA Taq polymerase; and 30 ng of DNA. The PCR cycles were: 94°C, for 3 min, followed by 30 cycles of 94°C for 1 min; annealing temperature from 55 to 57°C

for 1 min, in touchdown, 72°C, for 1 min, and final extension, at 72°C, for 5 min. Amplified products were separated through polyacrylamide gel electrophoresis.

Number and size of SSR amplification products were characterized. Qualitative analyses based on the presence or absence of SSR alleles were performed for characterization of polymorphism between segregating materials.

Results and Discussion

Thirty-five EST-SSR sequences were selected among 77 EST-SSR analyzed in 3,388 sequences expressed in different libraries from the EST Coffee Genome database (Table 1). In this database, the trinucleotides were the

Table 2. EST SSR primers designed from contigs available in the coffee EST database.

Locus	Contig	Primer sequence (5' – 3')	
LEG1	6239	F: GTG GCC AGT TGA GTT GCA TA	R: GAG ATG AGA TGG GCT TCT TGA
LEG2	846	F: GGA CAA AGA CGC CTA ATC CA	R: CCG TTG CAT GAT AAA CGA CA
LEG3	2664	F: GGA TAT CGC ATC AGC CTC TC	R: AGC GGT TTC TTT CTT TG
LEG4	2796	F: CCT TTG AGG TGG TCA GCC TA	R: AGA GAA CCC AGG GTC AAG TTT A
LEG5	4345	F: GTC CAT TAG GCA TTT GTT GC	G: CTG GCC GCT AAA GAT AAG TAG A
LEG6	7115	F: GGT TTA CCC CAT GAG AGC AG	R: CTT GTG CTT GAG CCA TTT CA
LEG7	8478	F: CGA GGA GTT GAT TGC TGG A	R: TTC GAG TGA AAT CCC AAT CC
LEG8	8911	F: ACG GAA GAG GCA GAG ACT GA	R: CTT CCT TGG ACC CTC ATC AA
LEG9	10226	F: AGG TTT CCA AAG GAG ATG AGC	R: GAA GAC AAG TCC ATC GTC CAA
LEG10	10393	F: CAT GAT CAA CCC AAC AGC AT	R: TGA ATG TGT ATC GGG AGC AG
LEG11	11764	F: CAC TGA AGG CCT GGA AGA AT	R: AGC ATC TGC AGC CTC CAT AG
LEG12	13870	F: CAC CAT AGC AAC TTC AAA CAC G	R: CAC ATC CAG GAA CCT TGC TC
LEG13	14035	F: GAA GAG GAA GAA GGG GCA AG	R: GTG GTG GAG GAA AGG GAT TC
LEG14	1174	F: AGA GCC CCT TCT CGC TAT GT	R: GTT AAC GCC GGA TAT GCT TC
LEG15	2291	F: TCT ATG CAC ACC CTC CAC AA	R: AGC ATT GGT TTG CTT CTT GC
LEG16	2913	F: CTG CTT GTT GGC CAT AAT GA	R: ACA GTG GAA GAA GGC GAG AG
LEG17	10221	F: TGC GAC ATT TCA CTC TGC AT	R: GCC TAT GTT GTG TAC GTG TGC
LEG18	1177	F: CAA ACA ATG GTG ACG TGC TC	R: CAA GGC CAG CGA AAA AGA T
LEG19	3388	F: GTT GCC GTC TTC AGC TCT TC	R: CAT TGG AAT TGT TGC TGG TG
LEG20	3982	F: ATC TGC AAC GAA AGC TGG AG	R: GGC TTT CCT TTG ACA ACT GG
LEG21	5374	F: GGA TGA TGG AAA AGC GAA TG	R: GGC CAA GTA GAG AGT GAT TTG C
LEG22	5376	F: CTT CAT CTC CCT GCC AAC AC	R: TCT TCT AAG GCC AGC AAG GA
LEG23	6480	F: ACA GCA GGG GAA GAG TGT GT	R: CCC AAT CTA AGC CCC TGA AT
LEG24	6708	F: CGT AGG CGA ACC AAA AAC TT	R: CAG GAC CCA AGG TCA CAT TC
LEG25	7368	F: CGA AGA CGA AGC ACC AAG AT	R: TAA TGG GTT CAA CCG GAG TC
LEG26	7797	F: TGA AGC TGC CTC CTG TTT CT	R: CGT CAG CTC AAG AAC TGT GC
LEG27	7861	F: ATA GAT TCC AGC GGA AAG CA	R: TGC TCC ACA CCA CTG CTA AC
LEG28	10553	F: TGT TCA CAG CTA AAC CCA ACC	R: TTG ACG GTG ACG ATG TTG AT
LEG29	11836	F: TGG CTT CCT TGA TGC TTT CT	R: TGT GAT GGA ACG CGA ATA TG
LEG30	12228	F: TTG CCA TGT TTC CCA TAC CT	R: CAA AGA TCA CCC CTC ATG CT
LEG31	13778	F: ACT ATT GCT CCC CCG GTA TC	R: GAC TCG GAG AAC ACG TTG GA
LEG32	13853	F: GGG TGA TGG AAA AGC AAA TG	R: CCA GCA TCA GCA AGT AAA AGG
LEG33	11356	F: TGT GAA AAA GCC TCG AGT GA	R: TAA CAC GAG GGA GGG AGA GA
LEG34	11478	F: GCA AAC TTG TGG CTG ATC G	R: ATC AAA CGC ACC CTG ACA TT
LEG35	11696	F: CCG ATG CTT CCC TAT TTC CT	R: CTC TTT GTG GCC ATC GAT CT

most abundant (39.02%), followed by di- (35.83%), tetra- (1.65%), hexa- (0.79%) and pentanucleotides (0.38%). The presence of mononucleotides was also significant (22.19%); however, these microsatellites are not greatly used since they have low polymorphism. In silico EST analyses performed on leaves and fruits of *C. canephora* (Poncet et al., 2004) showed that the trinucleotide microsatellites are also the most abundant, followed by the di- and hexanucleotides, representing 34.34, 25.75 and 22.04%, respectively. The significant presence of mononucleotides was also reported.

In comparison with some of the other approaches, EST-SSR may be slightly less polymorphic than genomic library-derived SSR, as there is pressure for sequence conservation in gene regions, reducing polymorphism (Scott, 2001). Selection of EST-SSRs with tri- and dinucleotide motifs was preferred to increase the probability of polymorphic EST markers being identified. Out of the 35 EST-SSR loci randomly selected, fragment amplification was detected in 27. Among these, eight loci were polymorphic, 18 were monomorphic, and, in the others, amplification was not observed or they showed diffuse bands (Table 1).

A total of 63 alleles were amplified and their sizes varied from 120 to 350 bp. These alleles showed amplified fragments with the expected size. The exception was LEG1 that amplified an allele with 600 bp that could be related to an intron. In most loci, two alleles were detected, but loci with three or four alleles were also found, which is expected for tetraploid species, such as *C. arabica*.

The LEG9, LEG11, LEG12, LEG13, LEG22, LEG26, LEG28 and LEG32 loci were polymorphic only among *C. racemosa* accessions and *C. arabica* progenies. Therefore, genetic variability is expected between these two species. Poncet et al. (2004) also observed EST-SSR polymorphism between

C. canephora for six species of *Coffea*; *C. eugenoides* S. Moore, *C. heterocalyx* Stoff., *C. liberica* W. Bull ex Hiern, *C. dewevrei* De Wild. & T. Durand, *Coffea* sp. Moloundou and *C. pseudozanguebariae* Bridson. The search for polymorphic SSR loci associated with a desirable trait is complex due to the limited number of coffee SSRs available. In species with a narrow genetic base, such as *C. arabica*, the probability of finding polymorphism is low (Baruah et al., 2003; Vieira et al., 2006). Genetic diversity among coffee species identified by EST-SSR markers can be used as a gene source in breeding programs, with potential for other studies of genetic mapping, in the identification of cultivars or in pre-breeding by the genotyping of germplasm bank accessions.

Polymorphism among the susceptible and resistant genotypes to the coffee leaf miner was observed in loci Sat 225 and Sat 229. The Sat 259 locus was monomorphic in all the evaluated genotypes. The amplification profile of the Sat 225 locus corresponds to a 300 bp allele (Figure 1). This allele was present in both susceptible and resistant bulks and also in *C. racemosa* accessions. As the *C. racemosa* parent no longer exists, a bulk of accessions from the IAC germplasm bank was used to represent the genetic diversity of this species. However, no polymorphic alleles associated with resistance to coffee leaf miner were detected. Another allele with approximately 270 bp was verified in the susceptible progenies (S4, S8, S15, S21, S24, S38, S42 and S48). For the Sat 229 primer, a 190 bp polymorphic allele was observed in the individuals of the S4, S8, S15, S21, S38 and S42 bulks.

After screening, bulks were opened and both Sat 225 and Sat 229 were evaluated in individuals from the F₅ Coromandel progenies. The Sat 225 amplification pattern was the same as that shown in




Figure 1. Amplification of the SSR Sat 225 locus in *Coffea racemosa* (CR1, CR2, CR3, CR4, CR6, CR8, CR9, CR10), bulks of resistant plants (R3, R23, R29, R30, R33, R34, R37, R43, R46, R49) and susceptible plants (S1, S4, S8, S13, S15, S21, S24, S38, S42, S48) to coffee leaf miner. The arrow indicates the 270 bp polymorphic band, found in the bulk of susceptible individuals

the bulks. The 270 bp allele, potentially associated with susceptibility to the coffee leaf miner, was present in 73% of the susceptible individuals and absent in 91% of the resistant individuals. Likewise, the 190 bp polymorphic allele from locus Sat 229 was observed in 70% of the susceptible individuals and absent in 94% of the resistant ones. According to Carvalho et al. (2008), a 30 to 40% of susceptibility to *L. coffeella* in 'Siriema' F₅ progenies is commonly observed in the field.

The F₆ population was characterized by the evaluation of lesions in the infested leaf disks. A total of 37 individuals were classified as resistant and 54 as susceptible to the insect. Molecular characterization shows that the 270 bp polymorphic allele from the Sat 225 locus was found in 91% of the susceptible genotypes and was absent in 36% of the resistant ones. The 190 bp polymorphic allele from locus Sat 229 was observed in 91% of the susceptible individuals and in 48% of the resistant ones. Although a high amplification frequency is observed in susceptible genotypes, both 270 bp (Sat 225) and 190 bp (Sat 229) alleles were observed in resistant genotypes. However, no significant correlation among the molecular markers and the presence of lesions in coffee leaves infested by *L. coffeella* was detected.

Herrera et al. (2009) evaluated the Sat 225 and Sat 229 loci in a F₂ segregating population resulting from a cross between the susceptible *C. arabica* cultivar Caturra and the resistant female parental hybrid derived from the cross between *C. arabica* and *C. canephora*. These authors observed the occurrence of a variable amplification pattern during the evaluated years, which may be associated with the infestation pressure of the pathogen. Genetic analysis indicates that the two loci could be associated with leaf-rust resistance and that the additive effect would be essential for the expression of resistance.

In the present work, alleles associated with resistance to the coffee leaf miner were not identified. The frequency of plants with resistance in segregating populations has not increased, remaining near 35%, although some families show a higher percentage. The genetic control of resistance to the coffee leaf miner appears to be more complex. Therefore, other factors, besides the control determined by the two genes, as mentioned by Medina Filho (1977) and Guerreiro Filho et al. (1999), may be involved.

The use of markers as tools to assist coffee breeding programs may contribute to save time for the development of new cultivars. However, the coffee molecular markers presently available have shown limited polymorphism. Several strategies for the development of new coffee molecular markers, such as diversity arrays technology (DARs) (Wittenberg et al., 2005) and single nucleotide polymorphism (SNPs), identified through comparative genomics or re-sequencing, using new generation sequencing, will certainly contribute to the development of more informative markers, which may be used to map valuable traits in a more precise manner.

Conclusions

1. SSR markers show inter- and intraspecific polymorphism in *Coffea arabica* and *C. racemosa*.
2. No polymorphic alleles are associated with resistance to coffee leaf miner in the evaluated SSR loci.

Acknowledgements

To Fundação Procafé and Instituto Agrônômico, for supplying the genetic material and for support in carrying out the research; to Consórcio Brasileiro de Pesquisa do Café, Conselho Nacional de Desenvolvimento Científico e Tecnológico and Fundação de Amparo à Pesquisa do Estado de Minas Gerais, for financial support.

References

- BARUAH, A.; NAIK, V.; HENDRE, P.S.; RAJKUMAR, R.; RAJENDRAKUMAR, P.; AGGARWAL, R.K. Isolation and characterization of nine microsatellite markers from *Coffea arabica* L., showing wide cross-species amplifications. **Molecular Ecology Notes**, v.3, p.647-650, 2003.
- CARVALHO, C.H.S. de; FAZUOLI, L.C.; GUERREIRO FILHO, O.; PEREIRA, A.A.; ALMEIDA, S.R. de; MATIELLO, J.B.; BARTHOLO, G.F.; SERA, T.; MOURA, W. de M.; MENDES, A.N.G.; REZENDE, J.C. de; FONSECA, A.F.A. da; FERRAO, M.A.G.; FERRÃO, R.G.; NACIF, A. de P.; SILVAROLLA, M.B.; BRAGHINI, M.T. Cultivares de café arábica de porte baixo. In: CARVALHO, C.H.S. de (Ed.). **Cultivares de café: origem, características e recomendações**. Brasília: Embrapa Café, 2008. p.157-226.
- DAVIS, A.P.; GOVAERTS, R.; BRIDSON, D.M.; STOFFELEN, P. An annotated taxonomic conspectus of the genus *Coffea* (Rubiaceae). **Botanical Journal of the Linnean Society**, v.152, p.465-512, 2006.

- FERREIRA, M.E.; GRATTAPAGLIA, D. **Introdução ao uso de marcadores moleculares em análise genética**. 2.ed. Brasília: Embrapa-Cenargen, 1996. 220p. (Embrapa-Cenargen. Documentos, 20).
- GUERREIRO FILHO, O. Coffee leaf miner resistance. **Brazilian Journal of Plant Physiology**, v.18, p.109-117, 2006.
- GUERREIRO FILHO, O.; SILVAROLLA, M.B.; ESQUES, A.B. Expression and mode of inheritance of resistance to leaf miner *Perileucoptera coffeella*. **Euphytica**, v.105, p.7-15, 1999.
- GUPTA, P.K.; VARSHNEY, R.K. The development and use of microsatellite markers for genetic analysis and plant breeding with emphasis on bread wheat. **Euphytica**, v.113, p.163-185, 2000.
- HERRERA, J.C.P.; ALVARADO, G.A.; CORTINA, H.A.G.; COMBES M.; ROMERO, G.G.; LASHERMES, P. Genetic analysis of partial resistance to coffee leaf rust (*Hemileia vastatrix* Berk & Br.) introgressed into the cultivated *Coffea arabica* L. from the diploid *C. canephora* species. **Euphytica**, v.167, p.57-67, 2009.
- LASHERMES, P.; COMBES, M.C.; ROBERT, J.; TROUSLOT, P.; D'HONT, A.; ANTHONY, F.; CHARRIER, A. Molecular characterisation and origin of the *Coffea arabica* L. genome. **Molecular and General Genetics**, v.261, p.259-266, 1999.
- MEDINA FILHO, H.P.; CARVALHO, A.; MEDINA, D.M. Germoplasma de *Coffea racemosa* e seu potencial de melhoramento do café. **Bragantia**, v.36, p.43-46, 1977.
- MONDEGO, J.M.C.; GUERREIRO FILHO, O.; BENGTON, M.H.; DRUMMOND, R.D.; FELIX, J.M.; DUARTE, M.P.; RAMIRO, D.A.; MALUF, M.P.; SOGAYAR, M.C.; MENOSSI, M. Isolation and characterization of *Coffea* genes induced during coffee leaf-miner (*Leucoptera coffeella*) infestation. **Plant Science**, v.169, p.351-360, 2005.
- MORGANTE, M.; HANAFEY, M.; POWELL, W. Microsatellites are preferentially associated with nonrepetitive DNA in plant genomes. **Nature Genetics**, v.30, p.194-200, 2002.
- PINTO, F. de O.; MALUF, M.P.; GUERREIRO FILHO, O. Study of simple sequence repeat markers from *Coffea* expressed sequences associated to leaf miner resistance. **Pesquisa Agropecuária Brasileira**, v.42, p.377-384, 2007.
- PONCET, V.; HAMON, P.; MINIER, J.; CARASCO, C.; HAMON, S.; NOIROT, M. SSR cross-amplification and variation within coffee trees (*Coffea* spp.). **Genome**, v.47, p.1071-1081, 2004.
- SCOTT, K.D. Microsatellite derived from ESTs, and their comparison with those derived by other methods. In: HENRY, R.J. (Ed.). **Plant genotyping: the DNA fingerprinting of plants**. Oxon: CABI, 2001. p.225-238.
- SOUZA, J.C. de; REIS, P.R.; RIGITANO, R.L. de O. **Bicho-mineiro do café: biologia, danos e manejo integrado**. 2.ed. Belo Horizonte: Epamig, 1998. 48p. (Epamig. Boletim técnico, 54).
- TAUTZ, D.; RENZ, M. Simple sequences are ubiquitous repetitive components of eukaryotic genomes. **Nucleic Acids Research**, v.12, p.4127-4138, 1984.
- VIEIRA, E.S.N.; ANDRADE, A.C.; COLOMBO, C.A. et al. Brazilian coffee genome project: an EST-based genomic resource. **Brazilian Journal of Plant Physiology**, v.18, p.95-108, 2006.
- WITTENBERG, A.H.J.; LEE, T. van der; CAYLA, C.; KILIAN, A.; VISSER, R.G.F.; SCHOUTEN, H.J. Validation of the high-throughput marker technology DArT using the model plant *Arabidopsis thaliana*. **Molecular Genetics and Genomics**, v.274, p.30-39, 2005.

Received on June 16, 2011 and accepted on November 9, 2011