

Notas Científicas

Somatic embryogenesis and plant regeneration in elite clones of *Theobroma cacao*

Thiago Édson Ribeiro da Silva⁽¹⁾, Luciana Cardoso Cidade⁽¹⁾, Fátima Cerqueira Alvim⁽¹⁾,
Júlio César de Mattos Cascardo⁽¹⁾ and Marcio Gilberto Cardoso Costa⁽¹⁾

⁽¹⁾Universidade Estadual de Santa Cruz, Centro de Biotecnologia e Genética, Departamento de Ciências Biológicas, Rodovia Ilhéus-Itabuna, Km 16, CEP 45662-000 Ilhéus, BA, Brazil. E-mail: thiagroe@gmail.com, lccidade@yahoo.com.br, alvim@uesc.br, cascardo@labbi.uesc.br, mcosta@labbi.uesc.br

Abstract – The objective of this work was to evaluate a procedure for somatic embryogenesis and regeneration of cacao (*Theobroma cacao* L.) elite clones. Petal explants from cacao clones TSH 565 and TSH 1188 were cultured on PCG and SCG-2 media, for calli growth. Somatic embryos were formed on the surface of embryogenic calli after transfer to embryo development (ED) medium. Clone TSH 565 showed a higher embryogenic potential than TSH 1188. The best combination of carbon source for embryo induction in ED medium was genotype-specific. Embryogenic callus formations increased in micropore tape-sealed Petri dishes, irrespective of cacao genotype. Mature somatic embryos were successfully converted into plantlets.

Index terms: biotechnology, cacao, gas exchange, micropropagation, morphogenesis.

Embrionese somática e regeneração in vitro de clones elite de *Theobroma cacao*

Resumo – O objetivo deste trabalho foi avaliar um procedimento para embrionese somática e regeneração de clones elite de cacau. Pétalas dos clones de cacau TSH 565 e TSH 1188 foram cultivadas em meios PCG e SCG-2 para o crescimento de calos. Embrões somáticos desenvolveram-se na superfície dos calos embrionéticos, após a transferência para o meio ED. O clone TSH 565 apresentou maior potencial embrionético do que o TSH 1188. A melhor combinação de fonte de carbono quanto à indução de embrões em meio ED foi específica do genótipo. A formação de calos embrionéticos foi superior em placas de Petri seladas com fita hipoalergênica, independentemente do genótipo. Embrões maduros de ambos os genótipos foram convertidos em plântulas.

Termos para indexação: biotecnologia, cacau, troca gasosa, micropropagação, morfogênese.

Theobroma cacao L. (cacao) is a tropical perennial tree, endemic to lowland rainforests of the Amazon Basin, and has been domesticated since pre-Columbian times (Hurst et al., 2002). Currently, cacao is grown throughout the humid tropics as a component of complex agroecosystems (Lobão, 2007) to supply the crescent global demand for cocoa, the processed form of cacao seeds used by the chocolate industry.

Since 1989, the witches-broom disease, caused by the hemibiotrophic fungus *Moniliophthora perniciosa* (= *Crinipellis perniciosa*), has spread throughout Brazil, destroying cacao tree (*Theobroma cacao* L.) plantations and leading to important economical and social changes in affected areas, such as the State of Bahia (Rocha et al., 1993; Purdy & Schmidt, 1996). The vegetative clonal propagation of elite cacao

genotypes, resistant to *M. perniciosa*, has been used as a potential means to recover cacao production in Brazil (Rudgard et al., 1993). However, there are some disadvantages associated with the current methods for vegetative propagation of cacao, via rooting or grafting of plagiotropic cuttings, including the intensive labor and associated costs, low propagation rate, and an undesirable bush-like growth pattern (Figueira & Janick, 1995).

The clonal propagation of cacao, through modern biotechnology techniques, can be greatly facilitated by multiplying elite materials using somatic embryogenesis. The main advantages of this method include the possibility of rapidly generating asexually propagated uniform plants of high genetic value, and the clonal production of orthotropic plants with

normal dimorphic architecture and taproot formation (Maximova et al., 2002). Somatic embryo production and plantlet regeneration have been achieved in a large number of genotypes (Li et al., 1998; Maximova et al., 2002), although practical utilization of this technology for clonal propagation remains hindered by an inability to successfully induce somatic embryogenesis in many elite cacao genotypes.

The objective of this work was to evaluate a procedure for somatic embryogenesis and plant regeneration from petal tissues of the elite cacao clones TSH 565 and TSH 1188, using the basic protocol described by Li et al. (1998) and Maximova et al. (2002). The effects of genotype, carbon source, and different Petri dish sealing material on somatic embryogenesis were also assessed.

Unopened immature flower buds about 4 to 5 mm long, from field-grown cacao plants, were collected early in the morning at Cepec/Ceplac (Centro de Pesquisas da Comissão Executiva do Plano da Lavoura Cacaueira, Ilhéus, Bahia, Brazil). Immature flower buds were surface-sterilized by immersion in 70% (v/v) ethanol, for 1 min and 30 s, followed by 20 min in a 2.5% (v/v) sodium hypochlorite solution containing 0.1% (v/v) Tween-20 and, then, five rinses in sterile distilled water. Explants were prepared by briefly blotting the immature flowers with sterile paper towels and slicing them perpendicular to their longitudinal axis at a position 1/3 of the flower length from the base, using a sterile scalpel blade. Petal tissues were extracted through the opening at the cut end using a sharp sterile forceps. Petal bases were used as explants source.

Petal base explants were first cultured on primary callus growth (PCG) medium (Li et al., 1998), with the abaxial surface in contact with the medium. Cultures were maintained in the dark at $27\pm 2^\circ\text{C}$ for 2 weeks, then transferred for another 2 weeks in a secondary callus growth (SCG-2) medium (Maximova et al., 2002). Somatic embryos were induced by transfer of floral tissue-derived calli from SCG-2 medium to embryo development (ED) medium (Maximova et al., 2002). Cultures were subcultured every 2 weeks in ED medium, under the same incubation conditions described above.

Various concentrations of sucrose, in combination or not with glucose in ED medium, were evaluated for their ability to stimulate somatic embryo production in the two cacao genotypes, as follows: ED4w/g–ED4w/g

(first subculture in ED medium, supplemented with 4% sucrose and 0.1% glucose – subsequent subcultures in same ED medium), ED4w/g–ED3w/g (first subculture in ED medium supplemented with 4% sucrose and 0.1% glucose – subsequent subcultures in ED medium supplemented with 3% sucrose and 0.1% glucose), ED3w/g–ED3w/g (first subculture in ED medium supplemented with 3% sucrose and 0.1% glucose – subsequent subcultures in same ED medium), ED4–ED4 (first subculture in ED medium supplemented with 4% sucrose – subsequent subcultures in same ED medium), ED4–ED3 (first subculture in ED medium supplemented with 4% sucrose – subsequent subcultures in ED medium supplemented with 3% sucrose), and ED3–ED3 (first subculture in ED medium supplemented with 3% sucrose – subsequent subcultures in same ED medium).

In order to evaluate the effects of Petri dish sealing material on frequency of somatic embryos production, petal base explants from both cacao genotypes were cultured in Petri dishes containing PCG, SCG-2, and ED (ED3w/g–ED3w/g) media, as describe above, sealed with two rounds of parafilm (American National, USA), polyvinyl chloride (PVC) transparent film, or hypoallergenic micropore tape (Micropore 3 M, Brazil).

Experiments were performed with 25 explants per Petri dish, five Petri dishes per treatment, and repeated once at least. The percentage of embryogenic and rhizogenic calli over the total number of cultured explants, representing the frequency of embryogenic and rhizogenic calli, respectively, and the average number of embryos per explant were determined two months after the culture initiation. Statistical analysis was performed with the software BIOESTAT (Universidade Federal do Pará, Brazil), which tested the experiments as a completely randomized design. Data of frequencies (%) were arc sen $x^{0.5}$ transformed prior to statistical analysis. Analysis of variance was applied and, for means comparison, Bonferroni's test was used, with a critical value at 5% probability.

The clone TSH 565 presented significantly higher frequency of embryogenic calli (42%) than the TSH 1188 (4.3%). The frequency of rhyzogenic calli in TSH 1188 was as twice as higher than the frequency of embryogenic calli, which indicates a need of further protocol improvement. The clone TSH 565 also produced significantly more embryos per explant (4.8) compared to TSH 1188 (1.6). However, the milky to

translucent rate of embryos was lower in TSH 565 (ca. 45%) than in TSH 1188 (ca. 70%). Milky embryos present higher conversion rates into plantlets, as compared to translucent embryos (Teixeira et al., 2002). Furthermore, they have cotyledons with higher embryogenic potential for secondary embryogenesis, increasing the efficiency and quality of embryos produced in vitro (Maximova et al., 2002).

Carbon source in ED medium significantly affected the rhizogenic calli frequency, the average number of embryos per explant, and the average number of milky embryos per explant produced by both cacao clones (Table 1). In general, rhizogenic calli were produced in higher frequencies, when explants from both clones were cultured on ED4–ED4 medium, although these differences were not significant for TSH 1188. A high number of embryos per explant was obtained in ED4w/g–ED4w/g and ED4w/g–ED3w/g media for TSH 565. For TSH 1188, more embryos per explant were produced in ED4w/g–ED3w/g, ED3w/g–ED3w/g,

and ED3–ED3 media. The clone TSH 565 produced significantly more milky embryos per explant, when cultured in ED4w/g–ED4w/g and ED4–ED3 media, while the best responses for TSH 1188 were obtained in ED4w/g–ED3w/g and ED3–ED3 media. Sucrose appears to be the most effective carbon source for somatic embryogenesis in general, and its combination with glucose (ED3w/g–ED3w/g) has been reported to be beneficial in cacao (Li et al., 1998). This study demonstrates that glucose is not considered to be an essential requirement for milky embryo production in TSH 565 and TSH 1188, since similar results can be obtained in the absence of this sugar, through variation or maintenance of sucrose levels in ED medium, depending on cacao genotype.

Petal explants cultured in Petri dishes, sealed with micropore tape produced significantly more embryogenic calli than those cultured in Petri dishes sealed with PVC or parafilm, irrespective of cacao genotype (Table 2). However, the differences were not

Table 1. Effect of different carbon sources on rhizogenic calli frequency, average number of embryos per explant, and average number of milky embryos per explant of petal explants, from TSH 565 and TSH 1188 cultured in embryo development (ED) medium⁽¹⁾.

Cacao clone	Carbon source (ED medium)	Rhizogenic calli frequency (%) ⁽²⁾	Average number of embryos per explant	Average number of milky embryos per explant
TSH 565	ED4w/g–ED4w/g	13b	5.53a	3.18a
	ED4w/g–ED3w/g	6b	5.55a	2.01ab
	ED3w/g–ED3w/g	13b	3.49c	1.39b
	ED4–ED4	26a	3.52c	1.36b
	ED4–ED3	8b	5.44ab	3.49a
	ED3–ED3	3b	4.74b	1.56b
TSH 1188	ED4w/g–ED4w/g	12 ^{ns}	0.04b	0.04b
	ED4w/g–ED3w/g	9	1.60a	1.60a
	ED3w/g–ED3w/g	10	1.03a	0.47b
	ED4–ED4	12	0.00c	0.00b
	ED4–ED3	6	1.00ab	0.20b
	ED3–ED3	9	1.60a	1.40a

⁽¹⁾Data from two independent experiments; means followed by the same letters, for each cacao clone, are not statistically different by Bonferroni's test, at 5% probability. ⁽²⁾Data of rhizogenic calli frequency (%) were transformed ($\text{arc sen } x^{0.5}$) prior to statistical analysis.

Table 2. Effect of different Petri dish sealing materials on embryogenic calli frequency and average number of embryos per explant of petal explants, from TSH 565 and TSH 1188 cultured in embryo development (ED) medium⁽¹⁾.

Cacao clone	Petri dish sealing material	Embryogenic calli frequency (%) ⁽²⁾	Average number of embryos per explant
TSH 565	PVC	19c	1.93 ^{ns}
	Parafilm	45b	1.80
	Micropore tape	65a	1.96
TSH 1188	PVC	2c	0.17 ^{ns}
	Parafilm	9b	0.12
	Micropore tape	13a	0.19

⁽¹⁾Data from two independent experiments; means followed by the same letters, for each cacao clone, are not statistically different by Bonferroni's test, at 5% probability. ⁽²⁾Data of embryogenic calli frequency (%) were transformed ($\text{arc sen } x^{0.5}$) prior to statistical analysis.

significant, when the average number of embryos and milky embryos per explant were evaluated. Carbon dioxide and ethylene are produced by all plant tissues cultured in vitro, they may accumulate to high levels inside culture vessels, under low light and reduced gas exchanges with the external environment, and may be detrimental or beneficial for cell growth and regeneration (Marino & Berardi, 2004). Micropore tape allows better gaseous exchange with the external environment, reducing carbon dioxide and ethylene accumulation inside the culture vessels.

Mature somatic embryos were selected for conversion into plantlets. It was observed that the conversion rate varied between the two cacao clones, and TSH 1188 showed higher rates than TSH 565. This observation could be related to morphological characteristics of the embryos. Besides more milky embryos, TSH 1188 presented higher rates of embryos with well-defined hypocotyl and cotyledons (normal embryos), while most of the embryos produced by TSH 565 presented fused hypocotyls and underdeveloped or extra cotyledons (abnormal embryos). About six months after culture initiation, plantlets of TSH 565 and TSH 1188, with more than three leaves and with healthy root systems, were successfully established in greenhouse.

Collectively, these data indicate that the cacao elite clones TSH 565 and TSH 1188 can be successfully regenerated from petal tissues, by using the somatic embryogenesis methods described by Li et al. (1998) and Maximova et al. (2002). Modifications of carbon source in ED medium and Petri dish sealing material improve the efficiency of the method. This procedure may allow the practical use of somatic embryogenesis for clonal propagation and genetic transformation of these important elite clones.

Acknowledgements

To Centro de Pesquisas da Comissão Executiva do Plano da Lavoura Cacaueira (Ilhéus, Bahia, Brazil), for providing the plant materials; to Conselho Nacional de Desenvolvimento Científico e Tecnológico and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, for fellowships granted; to International

Foundation for Science, for financial support; to Fundação de Amparo à Pesquisa do Estado da Bahia, for financial and fellowship support.

References

- FIGUEIRA, A.; JANICK, J. Somatic embryogenesis in cacao (*Theobroma cacao* L.). In: JAIN, S.; GUPTA, P.; NEWTON, R. (Ed.). **Somatic embryogenesis in woody plants**. Netherlands: Kluwer Academic Publisher, 1995. v.2. p.291-310.
- HURST, W.J.; TARKA, S.M.; POWIS, T.G.; VALDEZ, F.; HESTER, T.R. Cacao usage by the earliest Maya civilization. **Nature**, v.418, p.289-290, 2002.
- LI, Z.; TRAORE, A.; MAXIMOVA, S.; GUILTINAN, M.J. Somatic embryogenesis and plant regeneration from floral explants of cacao (*Theobroma cacao* L.) using thidiazuron. **In Vitro Cellular and Developmental Biology – Plant**, v.34, p.293-299, 1998.
- LOBÃO, D.E.V.P. **Agroecossistema Cacaueiro da Bahia: cacau-cabruca e fragmentos florestais na conservação de espécies arbóreas**. 2007. 98p. Tese (Doutorado) - Universidade Estadual Paulista Júlio de Mesquita Filho, Jaboticabal.
- MARINO, G.; BERARDI, G. Different sealing materials for Petri dishes strongly affect shoot regeneration and development from leaf explants of quince 'BA 29'. **In Vitro Cellular and Developmental Biology – Plant**, v.40, p.384-388, 2004.
- MAXIMOVA, S.N.; ALEMANNI, L.; YOUNG, A.; FERRIÈRE, N.; TRAORE, A.; GUILTINAN, M.J. Efficiency, genotypic variability, and cellular origin of primary and secondary somatic embryogenesis of *Theobroma cacao* L. **In Vitro Cellular and Developmental Biology – Plant**, v.38, p.252-259, 2002.
- PURDY, L.H.; SCHMIDT, R.A. Status of cocoa witches'-broom: biology, epidemiology, and management. **Annual Review of Phytopathology**, v.34, p.573-594, 1996.
- ROCHA, H.M.; MIRANDA, R.A.C.; SGRILLO, R.B.; SETUBAL, R.A. Witches'-broom in Bahia, Brazil. In: RUDGARD, S.A.; MADISON, A.C.; ANDEBRHAN, T. (Ed.). **Disease and management in cocoa: comparative epidemiology of witches'-broom**. London: Chapman and Hall, 1993. p.189-200.
- RUDGARD, S.A.; ANDEBRHAN, T.; MADDISON, A.C.; SCHMIDT, R.A. Disease management: recommendations. In: RUDGARD, S.A.; MADISON, A.C.; ANDEBRHAN, T. (Ed.). **Disease and management in cocoa: comparative epidemiology of witches'-broom**. London: Chapman and Hall, 1993. p.201-212.
- TEIXEIRA, J.B.; MARBACH, P.A.S.; SANTOS, M. de O. **Otimização da metodologia da embriogênese somática visando à propagação clonal de genótipos elite de cacau (*Theobroma cacao* L.)**. Brasília: Embrapa Recursos Genéticos e Biotecnologia, 2002. 34p. (Embrapa Recursos Genéticos e Biotecnologia. Documentos, 79).

Received on August 2, 2008 and accepted on September 25, 2008